New observations of the old magnetic nova GQ Muscae

W. Narloch[1], J. Kaluzny[1], W. Krzeminski[1], W. Pych[1], M. Rozyczka[1], S. Shectman[3], I. B. Thompson[3], T. Tomov[2]

[1] *Nicolaus Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland; wnarloch,jka,mnr,wk,pych@camk.edu.pl*

[2] *Torun Centre for Astronomy, Nicolaus Copernicus University, ul. Gagarina 11, 87-100, Torun, Poland; Toma.Tomov@astri.umk.pl*

[3] *Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101, USA; ian@ociw.edu*

arXiv:1401.7177v1 [astro-ph.SR] 28 Jan 2014

Abstract. We report photometric observations of GQ Mus performed between 1992 and 2011. We find that the total amplitude of the orbital modulation of its brightness decreased from ∼0.9 mag in 1992 to ∼0.2 mag in 2010, becoming comparable to the amplitude of chaotic flickering on a time scale of several minutes. Optical spectra obtained in 2001 and 2012 indicate continuing activity of GQ Mus. The spectra show broad emission lines of He II and H I typical for magnetic cataclysmic variables. The nova was observed as an UV-bright object in 2001 and 2012. We also show that the orbital period of GQ Mus has been constant between 1989 and 2010-2011.

Key words: stars: individual: GQ Mus – novae – cataclysmic variables

1. INTRODUCTION

The cataclysmic variable GQ Muscae (Nova Muscae 1983) is a classical novae. It was discovered by W. Liller on January 18th, 1983 (Shylaja 1983; Liller & Overbeek 1983). At the moment of discovery its brightness was estimated at 7.2 mag. The post-outburst optical light curve of GQ Mus was unusual: after an early decline lasting for about four months its brightness remained approximately constant for the next five months (Krautter et al. 1984). Duerbeck (1987) classified it as a moderately fast nova with $t_3 = 45$ d. With a pre-outburst magnitude of $V \geq 21$ (Krautter et al. 1984) GQ Mus belongs to a few known novae whose outburst amplitude approaches or even exceeds 14 mag.

Early spectra of GQ Mus were characterized by broad and strong HI and HeI emission lines. Lines of $FeII$, $TiII$ and $NIII$ were also visible, and they indicated a fast expansion of the ejected envelope (Charles 1983; Whitelock et al. 1983). Spectra obtained between 1984 and 1994 showed lines of highly ionized atoms, indicating a very high temperature of the central source of 2.65×10^5K (Diaz et al. 1995; Morisset & Pequignot 1996).

Photometric observations obtained in 1988 by Diaz & Steiner (1989) revealed a brightness modulation with a period of 85.5 min. The light curve was asymmetric, with the full range reaching 1 mag in the V-band. Further photometric and spectroscopic monitoring of allowed the classification of GQ Mus as a magnetic cataclysmic variable (Diaz & Steiner 1990, 1994) and that radial velocities measured from strong emission lines of $HeII\lambda4686$, $OIV\lambda3411$ and $NIII\lambda5270$ could be phased with the photometric period. The natural interpretation is that

Table 1. Summary of photometric observations of GQ Mus.

Detector	Season	No. of nights	No. of frames V	No. of frames B	Ave. exp. time in V[s]	<seeing> [arcsec]
TEK2	1992	4	165		185	1.5
SITe3	1997	3	21		122	1.3
	2000	1	21		300	3.0
	2001	3	43	2	300	1.5
	2002	6	173		300	1.8
TEK5	2008	1	5		60	1.5
SITe2k	2010	8	395	19	70	1.2
	2011	2	257	4	60	1.5

the 85.5 min periodicity corresponds to the orbital period of the system.

GQ Mus was the first classical nova with X-ray emission detected during an outburst (Ögelman et al. 1984). It remained in an active state for several years longer than predicted – supersoft X-ray emission persisted in 1992 (Ögelman et al. 1993). Later observations conducted in 1993 showed that the X-ray flux had decreased by a factor of 100 within one year (Shanley et al. 1994), indicating the extinction of the supersoft X-ray source.

These unusual properties of GQ Mus prompted us to embark on an occasional photometric monitoring of this object between 1992 and 2011. In the present paper we report the results of those observations. We also present two optical spectra collected in 2001 and 2012. The next section summarizes the observations, data reduction and calibrations. In section 3 we discuss the main results, and a brief summary of the paper is given in section 4.

2. OBSERVATIONS, DATA REDUCTION AND CALIBRATION

Photometric observations of GQ Mus were made with 1.0-m Swope and 2.5-m du Pont telescopes at Las Campanas Observatory in the period 1992–2011. The TEK2 and SITE3 CCD cameras were used on the Swope telescope, while the TEK5 and SITE2K cameras were employed on the du Pont telescope. A schematic log of the observations is given in Table 1. The variable was observed on 28 nights, yielding a total of 1080 V-band frames and 25 B-band frames. In 2001 we also obtained two frames in the$U$-band with SITE3/Swope.

Profile photometry was extracted using the Daophot/Allstar programs (Stetson 1987, 1990). The data from SITE3 were additionally reduced with the image subtraction code ISIS (Alard & Lupton 1998; Alard 2000). Measurements were transformed to the standard UBV sytem based on observations of ten standard stars from two Landolt fields (Landolt 1992) performed on June 17/18, 2001. The standard stars were observed at air-masses of 1.15 and 1.20 while GQ Mus was observed at an air mass of 1.27. Assuming average extinction coefficients for LCO, we obtained for GQ Mus $V = 18.65 \pm 0.02$ mag, $(B - V) = 0.27 \pm 0.02$ mag and $(U - B) = 0.78 \pm 0.03$ mag at $HJD = 2452078.48$. The data from the remaining cameras were transformed to the standard BV system based on local standards observed with SITE3. For most of the differential V photometry the accuracy of individual measurements ranges from 0.01 to 0.02 mag.

A medium resolution spectrum of the nova was obtained with the B%C spectrograph mounted on the Magellan Baade telescope. A 20 min. exposure centered

on 23h37m UT was made on June 18, 2001. The spectrum covered the range 3773-4490 Å at a resolution of 2800, and an average signal to noise ratio of 8. A second medium resolution spectrum of GQ Mus was collected with the South African Large Telescope on April 01, 2012. The RS spectrograph was used with the PC0900 grating and a 1.5 arcsec slit, providing a resolution of $R = 800$. Two 5 min exposures were centered on 0h:38m UT. The combined spectrum has a signal to noise ratio of 2.3 and 2.8 at $H\beta$ and $H\alpha$, respectively. The reduced and flux-calibrated spectrum covers the range 3800-6575 Å with 60-Å wide gaps at 4530 Å and 5590 Å.

3. RESULTS

In Fig. 1 we present a schematic V-light curve of GQ Mus based on data collected between 1992 and 2011. In that period the average luminosity of the nova dropped from <V> = 17.87 mag to <V> = 19.01 mag. Moreover, after 1992 the amplitude of the periodic variability significantly decreased. It is possible that around 2009 the luminosity of the nova stabilized, as no systematic change of <V> was observed during the subsequent two years. Note that in 2010-2011 the nova was still brighter than in the pre-outburst stage by at least two magnitudes.

We have analyzed yearly light curves for periodicity. For three seasons the amount of data was sufficient to establish the period with a high accuracy. We obtained $P = 0.05936538(\pm8)$ d, $P = 0.0593640(\pm2)$ d and $P = 0.05936553(\pm4)$ d for the years 1992, 2002 and 2010-2011, respectively, where the errors are 1-σ uncertainties of the last significant digit. Diaz and Steiner (1994) measured $P = 0.0593650(\pm1)$ d for 1989-1990. The combined measurements indicate that the orbital period of GQ Mus was constant between 1989 and 2011 at a level of $dP/P \leq 1.7 \times 10^{-6}$. Phased light curves for 1992, 2002 and 2010-2011 are presented in Fig. 2. The shape of the 1992 curve is generally similar to that obtained by Diaz and Steiner (1994) for the period 1989-1990. However, in 1992 the ascending branch was less steep, and the bump on the descending branch was less strongly pronounced. The light curves for 2002 and 2010-2011 have an amplitude of only ∼0.1 mag, and are noisy due to flickering on a time scale of a few minutes. The light curve for 1994 (Diaz et al. 1995) showed a significantly reduced amplitude in comparison with those for the 1989-1990 and 1992 seasons. Apparently the orbital modulation weakened with time, becoming comparable to the amplitude of flickering by the observing 2010-2011 seasons.

With galactic coordinates $l = 297.2$ deg and $b = -5.0$ deg, the nova resides in a rather crowded stellar field. For a heliocentric distance of $3.2 - 4.8$ kpc (Diaz & Steiner 1994) it is located 280-420 pc above the galactic disk, so that it must be either a thick disk or a halo object. Figure 3 shows $V/(B-V)$ and $V/(U-B)$ color-magnitude diagrams (CMD) of the 3×5 arcmin^2 field containing GQ Mus. The nova is the bluest object on both CMDs, and the only UV-bright star in the analyzed field. While the V-magnitude increased by about 0.4 mag between 2001 and 2010, the object became redder by just 0.06 mag.

Figure 4 shows the spectrum taken in June 2001, smoothed with a 7-pixel moving box. While smoothing reduces the resolution, it makes the broad Balmer emission lines more clearly visible. Such broad Balmer lines have been observed in several polars and intermediate polars in an active state (Rosen et al. 1993; Piirola et al. 2008; Thomas et al. 2012). The spectrum obtained with the SALT telescope on April 2012, smoothed with a 9-pixel moving box, is presented in Fig.

5. It was flux calibrated and then normalized at 5500 Å. Balmer lines together with $HeII\lambda4686$ and $NIII\lambda4650$ are seen in emission, with $H\alpha$ being particularly strong. The spectrum still exhibited signatures typical for magnetic cataclysmic variables in 2012. Moreover, the spectrum implies a very blue color for the nova (no correction for interstellar reddening has been made, so that GQ Mus was even brighter in the UV than the slope of the spectrum suggests).

4. CONCLUSIONS

Almost three decades after the outburst of GQ Mus the object was brighter than in the pre-outburst stage by at least two magnitudes. Its light curve exibited orbital modulation with a shape typical for magnetic CVs. In both 2001 and 2012 the nova was bright in the UV. Spectra obtained in 2001 and 2012 show broad emission lines of H and He, characteristic for polars. We found no evidence for period changes between 1989 and 2011 at an upper limit of $dP/P \leq 1.7 \times 10^{-6}$.

ACKNOWLEDGMENTS. This project was supported by grant DEC-2012/05/B/ST9/03931 from the Polish National Science Center.

REFERENCES

Alard C., Lupton R. H., 1998, ApJ, Vol. 503, p. 325
Alard C., 2000, A&A,
Charles P., 1983, IAU Circ., No. 3766
Diaz M. P., Steiner J. E., 1989, ApJ, 339, 41-43
Diaz M. P., Steiner J. E., 1990, Rev. Mex. Astron. Astrof., 21, 369-372
Diaz M. P., Steiner J. E., 1994, ApJ, 425, 252
Diaz M. P., Williams R. E., Phillips M. M., Hamuy M., 1995, MNRAS, 277, 959-964
Duerbeck H. W., 1987, SSRV, 45, 1-212
Krautter, J. et al., 1984, A&A, 137, 307
Landolt A. U., 1992, AJ, 104, nr 1, 340
Liller W., Overbeek M. D., 1983, IAU Circ., No. 3764
Morisset C., Pequignot D., 1996, A&A, 312, 135-159
Ögelman H., Beuermann K., Krautter J., 1984, ApJ, 287, 31-34
Ögelman H., Orio M., Krautter J., Starrfield S., 1993, Nature 361, 331
Piirola V., Vornanen T., Berdyugin A., Coyne S. J. G. V., 2008, ApJ, 684, 558
Rosen S. R., Mittaz J. P. D., Hakala P. J., 1993, MNRAS, 264, 171
Shanley L., Gallagher J. S., Ögelman H., Orio M., Krautter J., 1994, AAS, 184, 4601
Shylaja B. S., 1983, The Observatory, Vol. 103, p. 203-204
Stetson P. B., 1987, PASP, 99, 191-222
Stetson P. B., 1990, PASP, 102, 932-948
Thomas H. C., Beuermann K., Reinsch K., Schwope A. D., Burwitz, V., 2012, A&A, 546, 104
Whitelock P. A., Carter B. S., Feast M. W., Glass I. S., Laney D., Menzies J. W., Walsh J., Williams P. M., 1984, MNRAS, 211, 421-432



Fig. 1. The Light curve of GQ Mus for the period 1992-2011.



Fig. 2. Phased light curves of GQ Mus in V for seasons 1992, 2002 and 2010-2011.



Fig. 3. Color-magnitude diagrams for **the 3×5 arcmin^2 field centered on GQ Mus**. Positions of the variable are marked with triangles, with $V = \sim$18.5 mag corresponding to the season 2001, and $V = \sim$19.0 mag to the season 2010.



Fig. 4. The spectrum of GQ Mus from June 18, 2001 smoothed with a 7-pixel moving box.



Fig. 5. The spectrum of GQ Mus from April 2012 smoothed with a 9-pixel moving box.